

UCHL5IP Rabbit mAb [DNr5]

Cat NO. :A25365

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	Human	Q99871	37kDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:
Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human UCHL5IP

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Detected in spleen, thymus, testis, ovary, small intestine and colon, with highest levels of expression in testis and ovary..

Subcellular location:

 $\label{thm:cytoskeleton} \textbf{Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle.}$

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Contributes to mitotic spindle assembly, maintenance of centrosome integrity and completion of cytokinesis as part of the HAUS augmin-like complex..

Validation Data:

UCHL5IP Rabbit mAb [DNr5] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cell lysate. Using UCHL5IP Rabbit mAb [DNr5]at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com