gamma Synuclein Rabbit mAb [6yny]

Cat NO. :A28273

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC ICC IP FC	Human Rat	O76070	16kDa	Rabbit	lgG	50ul,100ul,200ul

Applications detail:

Application	Dilution		
WB	1:1000-2000		
ІНС	1:100		
The optimal dilutions should be	ne optimal dilutions should be determined by the end user		

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human gamma Synuclein

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol $\ (buffer)$.

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Highly expressed in brain, particularly in the substantia nigra. Also expressed in the corpus callosum, heart,

skeletal muscle, ovary, testis, colon and spleen. Weak expression in pancreas, kidney

Subcellular location:

Cytoplasm, perinuclear region. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm,

cytoskeleton, spindle.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Plays a role in neurofilament network integrity. May be involved in modulating axonal architecture during development and in the adult. In vitro, increases the susceptibility of neurofilament-H to calcium-dependent proteases (By similarity). May also function in modulating the keratin network in skin. Activates the MAPK and Elk-1 signal transduction pathway (By similarity)..

Validation Data:

gamma Synuclein Rabbit mAb [6yny] Images



Western blot (SDS PAGE) analysis of extracts from human fetal brain lysate.Using gamma Synuclein Rabbit mAb [6yny]at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.