

Guanylyl Cyclase beta 1 Rabbit mAb [qbv6]

Cat NO. :A37356

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC	Human,Mouse,R	Q02153	71kDa	Rabbit	IgG	50ul,100ul,200ul
	at					

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

_		•			
(:	าทเ	jug	ıatı	ρ.	
\sim		9	~~	•	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human Guanylyl Cyclase beta

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Detected in brain cortex and cerebellum (at protein level)...

Subcellular location:

Cytoplasm.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Mediates responses to nitric oxide (NO) by catalyzing the biosynthesis of the signaling molecule cGMP..

Validation Data:

Guanylyl Cyclase beta 1 Rabbit mAb [qbv6] Images



Western blot (SDS PAGE) analysis of extracts from Human fetal brain lysate. Using Guanylyl Cyclase beta 1 Rabbit mAb [qbv6]at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com