Cardiac Troponin C Rabbit mAb [Hi12]

Cat NO. :A39537

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC	Human,Mouse,R	P63316	18kDa	Rabbit	lgG	50ul,100ul,200ul
	at					

Applications detail:

Application	Dilution		
WB	1:1000-2000		
ІНС	1:100		
The optimal dilutions should be determined by the end user			

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human Cardiac Troponin C

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Troponin is the central regulatory protein of striated muscle contraction. Tn consists of three components: Tn-I which is the inhibitor of actomyosin ATPase, Tn-T which contains the binding site for tropomyosin and Tn-C. The binding of calcium to Tn-C abolishes the inhibitory action of Tn on actin filaments.

Validation Data:

Cardiac Troponin C Rabbit mAb [Hi12] Images



Western blot (SDS PAGE) analysis of extracts from Human fetal heart lysate.Using Cardiac Troponin C Rabbit mAb [Hi12]at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.