

P-PKC delta (T507) Rabbit mAb [ZJ1f]

Cat NO. :A41256

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	Human	Q05655(PKC delta, 抗	78kDa	Rabbit	IgG	50ul,100ul,200ul
		原)P05129(PKC gamma),				
		P05771(PKC beta),				
		P17252(PKC alpha),				
		Q02156(PKC epsilon),				
		Q05513(PKC zeta)				

Applications detail:	Application	Dilution	
	WB	1:1000-2000	
	The optimal dilutions should be determined by the end user		

Conjugate:
UnConjugate
Form:
Liquid
sensitivity:
Endogenous
Purification:
Affinity-chromatography
Specificity:
Antibody is produced by immunizing animals with A synthesized peptide derived from human Phospho-PKC delta
(T507)

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



• • •					•
บวเ		12ti	n	110	ıta:
va	пч	аы	UII	UC	ıLa:

P-PKC delta (T507) Rabbit mAb [ZJ1f] Images

Western blot (SDS PAGE) analysis of extracts from 0.Using P-PKC delta (T507) Rabbit mAb [ZJ1f]at dilution of 1:1000 incubated at 4 $^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.