

# MUC5AC Rabbit mAb [9V1R]

Cat NO. :A42452

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size	
WB IHC ICC/IF	Human	P98088	220-600kDa	Rabbit	IgG	50ul,100ul,200ul	

Applications detail:	Application	Dilution
	WB	1:1000-2000
	ІНС	1:100
	ICC/IF	1:100
	The optimal dilutions should be de	etermined by the end user

				te:		
	$\sim$ 10		~	~	-	-
$\mathbf{u}$	UH	IIU	u	а	LE	=

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

#### Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human MUC5AC

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

## Tissue specificity:

Highly expressed in surface mucosal cells of respiratory tract and stomach epithelia. Overexpressed in a number of carcinomas. Also expressed in Barrett's esophagus epithelium and in the proximal

## Subcellular location:

Secreted.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Gel-forming glycoprotein of gastric and respiratory tract epithelia that protects the mucosa from infection and chemical damage by binding to inhaled microorganisms and particles that are subsequently removed by the mucociliary system (PubMed:14535999, PubMed:14718370). Interacts with H.pylori in the gastric epithelium, Barrett's esophagus as well as in gastric metaplasia of the duodenum (GMD) (PubMed:14535999)..

## **Validation Data:**

## MUC5AC Rabbit mAb [9V1R] Images



View more information on http://naturebios.com