

# KAT8 / MYST1 / MOF Rabbit mAb [1DzQ]

Cat NO. :A58493

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC ICC/IF IP	Human,Mouse,R	Q9H7Z6	52kDa	Rabbit	IgG	50ul,100ul,200ul
FC	at					

Applications detail:

Application

WB

1:1000-2000

IHC

1:100

ICC/IF

1:100

The optimal dilutions should be determined by the end user

$\sim$		•			
( : (	nnı	חווו	ate	٠.	
•	<b>UI I</b>	ч	acc	•	

UnConjugate

Form:

Liquid

sensitivity:

**Endogenous** 

Purification:

Affinity-chromatography

## Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human KAT8 / MYST1 / MOF

# Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

### Subcellular location:

Nucleus. Chromosome.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Histone acetyltransferase which may be involved in transcriptional activation (PubMed:12397079, PubMed:22020126). May influence the function of ATM (PubMed:15923642). As part of the MSL complex it is involved in acetylation of nucleosomal histone H4 producing specifically H4K16ac (PubMed:16227571, PubMed:16543150, PubMed:21217699, PubMed:22547026, PubMed:22020126). As part of the NSL complex it may be involved in acetylation of nucleosomal histone H4 on several lysine residues (PubMed:20018852, PubMed:22547026). That activity is less specific than the one of the MSL complex (PubMed:20018852, PubMed:22547026). Can also acetylate TP53/p53 at 'Lys-120'..

### **Validation Data:**

#### KAT8 / MYST1 / MOF Rabbit mAb [1DzQ] Images



Western blot ( SDS PAGE ) analysis of extracts from HeLa cell lysate. Using KAT8 / MYST1 / MOF Rabbit mAb [1DzQ]at dilution of 1:1000 incubated at  $4\,^\circ\!\mathrm{C}$  over night.

View more information on http://naturebios.com