

RASA1 Rabbit mAb [5QQh]

Cat NO. :A63604

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC	Human Rat	P20936	140kDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:	Application	Dilution
	wв	1:1000-2000
	ІНС	1:100
	The optimal dilutions should be d	etermined by the end user

\sim		•		
	n		Oto	
\sim	<i>)</i>	uu	ate	-

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:
Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human RASA1

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

In placental villi, detected only in the trophoblast layer (cytotrophoblast and syncytiotrophoblast). Not detected in stromal, endothelial or Hofbauer cells (at protein level)..

Subcellular location:

Cytoplasm.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Inhibitory regulator of the Ras-cyclic AMP pathway. Stimulates the GTPase of normal but not oncogenic Ras p21, this stimulation may be further increased in the presence of NCK1..

Validation Data:

RASA1 Rabbit mAb [5QQh] Images



Western blot (SDS PAGE) analysis of extracts from Rat brain cell lysate. Using RASA1 Rabbit mAb [5QQh]at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com