

NPRC Rabbit mAb [80Km]

Cat NO. :A68710

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB FC	Human,Mouse,R	P17342	70kDa	Rabbit	IgG	50ul,100ul,200ul
	at					

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

_						
\mathbf{C}	าทเ	LIC	gat	e:		

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human Natriuretic Peptide Receptor C

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Membrane, Single-pass type I membrane protein.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Receptor for the natriuretic peptide hormones, binding with similar affinities atrial natriuretic peptide NPPA/ANP, brain natriuretic peptide NPPB/BNP, and C-type natriuretic peptide NPPC/CNP. May function as a clearance receptor for NPPA, NPPB and NPPC, regulating their local concentrations and effects. May regulate diuresis, blood pressure and skeletal development. Does not have guanylate cyclase activity..

Validation Data:

NPRC Rabbit mAb [80Km] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cell lysate. Using NPRC Rabbit mAb [80Km]at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com