

TTR Rabbit mAb [9IS8]

Cat NO. :A70861

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC ICC/IF IP	Human	P02766	16kDa	Rabbit	IgG	50ul,100ul,200ul
FC						

Applications detail:

Application Dilution

WB 1:1000-2000

IHC 1:100

ICC/IF 1:100

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human TTR

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Detected in serum and cerebrospinal fluid (at protein level). Highly expressed in choroid plexus epithelial cells.

Detected in retina pigment epithelium and liver..

Subcellular location:

Secreted. Cytoplasm.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Thyroid hormone-binding protein. Probably transports thyroxine from the bloodstream to the brain..

Validation Data:

TTR Rabbit mAb [9IS8] Images



Western blot (SDS PAGE) analysis of extracts from human serum lysate. Using TTR Rabbit mAb [9IS8]at dilution of 1:1000 incubated at $4\,^\circ\!\mathrm{C}$ over night.

View more information on http://naturebios.com