ULBP1 Rabbit mAb [8Hgv]

Cat NO. :A79162

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	Human	Q9BZM6	36kDa	Rabbit	lgG	50ul,100ul,200ul

Applications detail:

Application Dilution WB 1:1000-2000 The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human ULBP1

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in T-cells, B-cells, erythroleukemia cell lines and in a wide range of tissues including heart, brain, lung,

liver, testis, lymph node, thymus, tonsil and bone marrow. Also found in fetal

Subcellular location:

Cell membrane, Lipid-anchor, GPI-anchor. Endoplasmic reticulum.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Binds and activates the KLRK1/NKG2D receptor, mediating natural killer cell cytotoxicity..

Validation Data:

ULBP1 Rabbit mAb [8Hgv] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cell lysate. Using ULBP1 Rabbit mAb [8Hgv]at dilution of 1:1000 incubated at 4 $^\circ\!C$ over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.