

NPHS2 Rabbit mAb [aM8c]

Cat NO. :A86053

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC	Human,Mouse,R	Q9NP85	42kDa	Rabbit	IgG	50ul,100ul,200ul
	at					

Applications detail: Application WB

Dilution 1:1000-2000

IHC

1:100

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human NPHS2

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli.

Subcellular location:

[Isoform 1]: Cell membrane,Peripheral membrane protein.,[Isoform 2]: Endoplasmic reticulum.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.

Validation Data:

NPHS2 Rabbit mAb [aM8c] Images



Western blot (SDS PAGE) analysis of extracts from Human fetal kidney lysate. Using NPHS2 Rabbit mAb [aM8c]at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com