

HOXA5 Rabbit mAb [k2WC]

Cat NO. :A97309

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	Human,Mouse,R	P20719	43kDa	Rabbit	IgG	50ul,100ul,200ul
	at					

****	Truman, wouse, ix	1 207 13	TORDA	Nabbit	igo	3001,10001,20			
	at								
Applications detail:		Application	Application Dilution						
		WB				1:1000-2000			
		The optimal	The optimal dilutions should be determined by the end user						
Conjugate:									
UnConjugate									
Form:									
Liquid									
sensitivity:									
Endogenous									
Purification:									
Affinity-chromato	graphy								
Specificity:									
Antibody is produ	ced by immuniz	ing animals with A sy	nthesized pepti	de derived	l from humar	1 HOXA5			
Storage buff	er and cond	ditions:							
Antibody store in	10 mM PBS, 0.5	mg/ml BSA, 50% glyc	erol (buffer) .						
Shipped at 4°C. S	tore at-20°C or -	-80°C.							
Products are valid	d for one natura	l year of receipt.Avoid	d repeated free:	ze / thaw c	ycles.				
Tissue speci	ficity:								
Subcellular I	ocation:								
Nucleus.									
Function:									

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'.

Validation Data:

HOXA5 Rabbit mAb [k2WC] Images



Western blot (SDS PAGE) analysis of extracts from 293T cell lysate. Using HOXA5 Rabbit mAb [k2WC]at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com