

# AceCS1 Rabbit mAb [A3L8]

Cat NO. :A11838

### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	Q9NR19	78 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

					te:		
	$\sim$ n		$\sim$	$\sim$	_	-	
$\mathbf{u}$	OH	IIU	u	а		=	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

## Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of Human AceCS1

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

## Tissue specificity:

## Subcellular location:

Cytoplasm, cytosol.

#### Function:

Catalyzes the synthesis of acetyl-CoA from short-chain fatty acids (PubMed:10843999, PubMed:28003429). Acetate is the preferred substrate (PubMed:10843999, PubMed:28003429). Can also utilize propionate with a much lower affinity (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



# **Validation Data:**

# AceCS1 Rabbit mAb [A3L8] Images



Western blot (SDS PAGE) analysis of extracts from LN18 cells.Using AceCS1 Rabbit mAb [A3L8] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com