

BCAT2 Rabbit mAb [VV3F]

Cat NO. :A18268

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	O15382	39 kDa	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution
	WB	1:1000-2000
	The optimal dilutions should b	e determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of Human BCAT2

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Ubiquitous..

Subcellular location:

Mitochondrion.

Function:

Catalyzes the first reaction in the catabolism of the essential branched chain amino acids leucine, isoleucine, and valine (PubMed:8702755, PubMed:25653144, PubMed:17050531). May also function as a transporter of branched chain alpha-keto acids (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

BCAT2 Rabbit mAb [VV3F] Images



Western blot (SDS PAGE) analysis of extracts from MCF-7 cells.Using BCAT2 Rabbit mAb [VV3F] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com