

RAE1 Rabbit mAb [679E]

Cat NO. :A19469

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	н	P78406	41 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application Dilution
WB 1:1000-2000
IHC 1:100,

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human RAE1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

 $\label{thm:cytoplasm} \textbf{Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, spindle pole. Nucleus envelope.}$

Function:

Acts as mRNA export factor involved in nucleocytoplasmic transport (PubMed:33849972, PubMed:20498086). Plays a role in mitotic bipolar spindle formation (PubMed:17172455). May function in attaching cytoplasmic mRNPs to the cytoskeleton both directly or indirectly (PubMed:17172455)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

RAE1 Rabbit mAb [679E] Images



Western blot (SDS PAGE) analysis of extracts from MCF-7 cells lyastes.using RAE1 Rabbit mAb [679E] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night

View more information on http://naturebios.com