

Caspase-7 Rabbit mAb [NB8E]

Cat NO. :A24429

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	P55210	32,35 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

\sim		•	gate:			
1.0	n		Oto			
\sim	<i>)</i>	uu	ıaıc	-		

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human Caspase-7

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Highly expressed in lung, skeletal muscle, liver, kidney, spleen and heart, and moderately in testis. No expression in the brain.

Subcellular location:

Cytoplasm.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Involved in the activation cascade of caspases responsible for apoptosis execution (PubMed:8521391, PubMed:8576161, PubMed:8567622, PubMed:9070923, PubMed:11701129). Cleaves and activates sterol regulatory element binding proteins (SREBPs) (PubMed:8521391, PubMed:8576161, PubMed:8567622, PubMed:9070923). Proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-|-Gly-217' bond. Overexpression promotes programmed cell death (PubMed:8521391, PubMed:8576161, PubMed:8567622, PubMed:9070923). Cleaves phospholipid scramblase proteins XKR4, XKR8 and XKR9 (By similarity)..., [Isoform Beta]: Lacks enzymatic activity...

Validation Data:

Caspase-7 Rabbit mAb [NB8E] Images



Western blot (SDS PAGE) analysis of extracts from wild-type HAP1 cells.Using Caspase-7Rabbit mAb [NB8E] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com