SCGN/Secretagogin Rabbit mAb [3894]

Cat NO. :A25699

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,R	O76038	32 kDa	Rabbit	lgG	100ul,200ul

Applications detail:

Application	Dilution			
WB	1:1000-2000			
IHC	1:100,			
The optimal dilutions should be d	e optimal dilutions should be determined by the end user			

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human SCGN/Secretagogin.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed at high levels in the pancreatic islets of Langerhans and to a much lesser extent in the gastrointestinal

tract (stomach, small intestine and colon), the adrenal medulla and cortex and the

Subcellular location:

Cytoplasm. Secreted. Cytoplasmic vesicle, secretory vesicle membrane,Peripheral membrane protein,Cytoplasmic side.

Function:


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

SCGN/Secretagogin Rabbit mAb [3894] Images



Western blot (SDS PAGE) analysis of extracts from Fetal brain tissue lyaste.using SCGN/Secretagogin Rabbit mAb [3894] at dilution of 1:1000 incubated at 4°C over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.