

DAZAP1 Rabbit mAb [3W8C]

Cat NO. :A26889

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	Q96EP5	43 kda	Rabbit	IgG	50ul,100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

\sim	•					
1''	าทแ	IMATA				
\mathbf{v}	JIIIL	ıgate	•			

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human DAZAP1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Mainly expressed in testis. Expressed to a lower level in thymus. Weakly or not expressed in heart, liver, brain, placenta, lung, skeletal muscle, kidney and pancreas..

Subcellular location:

Cytoplasm. Nucleus.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



RNA-binding protein, which may be required during spermatogenesis.

Validation Data:

DAZAP1 Rabbit mAb [3W8C] Images



Western blot (SDS PAGE) analysis of extracts from Jurkat cells lyastes.using DAZAP1 Rabbit mAb [3W8C] at dilution of 1:1000 incubated at $4\,^{\circ}\mathrm{C}$ over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.