

BCL2L14/BCLG Rabbit mAb[3C7X]

Cat NO. :A28266

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	Q9BZR8	37KDa	Rabbit	IgG	50ul 100ul,200ul

Applications detail: A

Application Dilution
WB 1:1000-2000
IHC 1:100

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human BCL2L14/BCLG.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Isoform 1 is widely expressed. Isoform 2 is testis-specific..

Subcellular location:

Cytoplasm.,[Isoform 1]: Cytoplasm, cytosol.

Function:

Plays a role in apoptosis.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

BCL2L14/BCLG Rabbit mAb[3C7X] Images



Western blot (SDS PAGE) analysis of extracts from A-549 cells.Using BCL2L14/BCLG Rabbit mAb IgG [3C7X] at dilution of 1:1000 incubated at $4\,^\circ\!\mathrm{C}$ over night.

View more information on http://naturebios.com