

# TPMT Rabbit mAb[B87C]

Cat NO. :A29260

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	P51580	32KDa	Rabbit	IgG	50ul 100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human TPMT.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

Tissue specificity:

### Subcellular location:

Cytoplasm.

### Function:

Catalyzes the S-methylation of thiopurine drugs such as 6-mercaptopurine (also called mercaptopurine, 6-MP or its brand name Purinethol) and 6-thioguanine (also called tioguanine or 6-TG) using S-adenosyl-L-methionine as the methyl donor (PubMed:657528, PubMed:18484748). TPMT activity modulates the cytotoxic effects of thiopurine prodrugs. A natural substrate for this enzyme has yet to be identified..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## TPMT Rabbit mAb[B87C] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells.Using TPMT Rabbit mAb IgG [B87C] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com