# HK3 Mouse mAb[I02B]

Cat NO. :A29738

## Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	P52790	99kDa	Mouse	lgG	50ul 100ul,200ul

## **Applications detail:**

Application	Dilution		
WB	1:1000-2000		
ІНС	1:100		
The optimal dilutions should be determined by the end user			

## Conjugate:

UnConjugate

Form:

Liquid

## sensitivity:

Endogenous

# **Purification**:

Protein A purification

#### Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human HK3.

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

**Tissue specificity**:

## Subcellular location:

### **Function**:

Catalyzes the phosphorylation of hexose, such as D-glucose and D-fructose, to hexose 6-phosphate (D-glucose 6-phosphate and D-fructose 6-phosphate, respectively) (PubMed:8717435). Mediates the initial step of glycolysis by catalyzing phosphorylation of D-glucose to D-glucose 6-phosphate (PubMed:8717435).


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

# Validation Data:

# HK3 Mouse mAb[I02B] Images



Western blot (SDS PAGE) analysis of extracts from Raji cells.Using HK3 Mouse mAb IgG [I02B] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.

For Research Use Only. Not For Use In Diagnostic Procedures.