

D4 GDI Rabbit mAb [NE61]

Cat NO. :A32362

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	P52566	23 kDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:

Application

WB 1:1000-2000

IHC 1:100,

ICC/IF 1:100,

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human D4 GDI.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Detected in bone marrow, thymus and spleen..

Subcellular location:

Cytoplasm, cytosol.

Function:

Regulates the GDP/GTP exchange reaction of the Rho proteins by inhibiting the dissociation of GDP from them, and the subsequent binding of GTP to them (PubMed:8356058, PubMed:7512369). Regulates reorganization of the actin cytoskeleton mediated by Rho family members (PubMed:8262133)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

D4 GDI Rabbit mAb [NE61] Images



Western blot (SDS PAGE) analysis of extracts from Jurkat cells lyastes.using D4 GDI Rabbit mAb [NE61] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night

View more information on http://naturebios.com