

# PCK2 Rabbit mAb [8BV9]

Cat NO. :A33833

### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	Н,М	Q16822	71 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

$\sim$	•					
1''	าทแ	IMATA				
$\mathbf{v}$	JIIIL	ıgate	•			

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human PCK2.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

## Subcellular location:

Mitochondrion.

#### Function:

Catalyzes the conversion of oxaloacetate (OAA) to phosphoenolpyruvate (PEP), the rate-limiting step in the metabolic pathway that produces glucose from lactate and other precursors derived from the citric acid cycle...

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## PCK2 Rabbit mAb [8BV9] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells lyastes.using PCK2 Rabbit mAb [8BV9] at dilution of 1:1000 incubated at  $4\,^\circ\!\mathrm{C}$  over night

View more information on http://naturebios.com