## MC-2 Rabbit mAb [A915]

Cat NO. :A33943

## Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	н	P33032	45 kDa	Rabbit	lgG	100ul,200ul

### **Applications detail:**

Application	Dilution	
WB	1:1000-2000	
ІНС	1:100,	
he optimal dilutions should be determined by the end user		

### Conjugate:

UnConjugate

Form:

Liquid

#### sensitivity:

Endogenous

## **Purification**:

Protein A purification

#### Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human MC-2.

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

#### **Tissue specificity:**

Expressed in the brain but not in the melanoma cells.

## Subcellular location:

Cell membrane, Multi-pass membrane protein.

#### **Function**:

Receptor for MSH (alpha, beta and gamma) and ACTH. The activity of this receptor is mediated by G proteins which activate adenylate cyclase. This receptor is a possible mediator of the immunomodulation properties of melanocortins.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

# Validation Data:

## MC-2 Rabbit mAb [A915] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells lyastes.using MC-2 Rabbit mAb [A915] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.