

Prostein Mouse mAb[7066]

Cat NO. :A34013

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size	
WB,IHC,ICC/IF	н	Q96JT2	59kda	Mouse	IgG	100ul,200ul	

Applications detail:

ApplicationDilutionWB1:1000-2000IHC1:100ICC/IF1:100The optimal dilutions should be determined by the end user

	onjug							
,	\sim	n		\sim	$\boldsymbol{\circ}$	ナハ		
.,	.,				~		=	
_	_	• • •	_		•		-	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human Prostein.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Prostate specific. Expressed in all prostatic glandular cells. Expressed both in normal and cancerous prostates..

Subcellular location:

 ${\bf Membrane, Multi-pass\ membrane\ protein.}$

Function:


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

Prostein Mouse mAb[7066] Images



Western blot (SDS PAGE) analysis of extracts from rat testis tissue. Using Prostein Mouse mAb IgG [7066] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com