ABHD5 Mouse mAb[2R57]

Cat NO. :A37239

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	н	Q8WTS1	39kDa	Mouse	lgG	100ul,200ul

Applications detail:

Application	Dilution	
WB	1:1000-2000	
ICC/IF	1:100	
The optimal dilutions should be determined by the end user		

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human ABHD5.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Widely expressed in various tissues, including lymphocytes, liver, skeletal muscle and brain. Expressed by upper

epidermal layers and dermal fibroblasts in skin, hepatocytes and neurons (at protein

Subcellular location:

Cytoplasm. Lipid droplet. Cytoplasm, cytosol.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Coenzyme A-dependent lysophosphatidic acid acyltransferase that catalyzes the transfert of an acyl group on a lysophosphatidic acid (PubMed:18606822). Functions preferentially with 1-oleoyl-lysophosphatidic acid followed by 1-palmitoyl-lysophosphatidic acid, 1-stearoyl-lysophosphatidic acid and 1-arachidonoyl-lysophosphatidic acid as lipid acceptor. Functions preferentially with arachidonoyl-CoA followed by oleoyl-CoA as acyl group donors (By similarity). Functions in phosphatidic acid biosynthesis (PubMed:18606822). May regulate the cellular storage of triacylglycerol through activation of the phospholipase PNPLA2 (PubMed:16679289). Involved in keratinocyte differentiation (PubMed:18832586). Regulates lipid droplet fusion (By similarity)...

Validation Data:

ABHD5 Mouse mAb[2R57] Images



Western blot (SDS PAGE) analysis of extracts from LNCaP cells.Using ABHD5 Mouse mAb IgG [2R57] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.