

RECQ1 Rabbit mAb [IR0U]

Cat NO. :A38173

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	P46063	72 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB 1:1000-2000

IHC 1:100,

ICC/IF 1:100,

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human RECQ1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

 $\label{thm:light} \mbox{High expression in heart, lung, skeletal muscle and kidney, low expression in brain.}$

Subcellular location:

Nucleus.

Function:

DNA helicase that may play a role in the repair of DNA that is damaged by ultraviolet light or other mutagens.

Exhibits a magnesium-dependent ATP-dependent DNA-helicase activity that unwinds single- and double-stranded DNA in a 3'-5' direction..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

RECQ1 Rabbit mAb [IR0U] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells lyastes.using RECQ1 Rabbit mAb [IR0U] at dilution of 1:1000 incubated at $4\,^\circ\mathrm{C}$ over night

View more information on http://naturebios.com