

Angiopoietin 2 Rabbit mAb [6MDG]

Cat NO. :A44285

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	O15123	57 kDa	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution
	WB	1:1000-2000
	The optimal dilutions should be	determined by the end user

\sim		•	gate:		
	\mathbf{n}		ハナハ	-	
v	UH	uu	ıaıe	-	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human Angiopoietin 2.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Secreted.

Function:

Binds to TEK/TIE2, competing for the ANGPT1 binding site, and modulating ANGPT1 signaling (PubMed:15284220, PubMed:19116766, PubMed:19223473,PubMed:9204896). Can induce tyrosine phosphorylation of TEK/TIE2 in the absence of ANGPT1 (PubMed:15284220, PubMed:19116766, PubMed:19223473,PubMed:9204896). In the absence of angiogenic inducers, such as VEGF, ANGPT2-mediated loosening of cell-matrix contacts may induce endothelial cell apoptosis with consequent vascular regression. In concert with VEGF, it may facilitate endothelial cell migration and proliferation, thus serving as a permissive

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



angiogenic signal (PubMed:15284220, PubMed:19116766, PubMed:19223473, PubMed:9204896). Involved in the regulation of lymphangiogenesis (PubMed:32908006)..

Validation Data:

Angiopoietin 2 Rabbit mAb [6MDG] Images



Western blot (SDS PAGE) analysis of extracts from Mouse small intestine tissue lyaste.using Angiopoietin 2 Rabbit mAb [6MDG] at dilution of 1:1000 incubated at

View more information on http://naturebios.com