

Ferritin Rabbit mAb [4VJW]

Cat NO. :A46681

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	H,M,R	P02794	19,21 kDa	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution
	WB	1:1000-2000
	ICC/IF	1:100
	The optimal dilutions should be	determined by the end user

\sim	•		•
1''	าทแ	100	to.
\mathbf{v}	JIII	uga	LE:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human Ferritin

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in the liver..

Subcellular location:

Function:

Stores iron in a soluble, non-toxic, readily available form. Important for iron homeostasis. Has ferroxidase activity. Iron is taken up in the ferrous form and deposited as ferric hydroxides after oxidation. Also plays a role in delivery of iron to cells. Mediates iron uptake in capsule cells of the developing kidney (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

Ferritin Rabbit mAb [4VJW] Images



Western blot (SDS PAGE) analysis of extracts from HEK293T cells.Using FerritinRabbit mAb [4VJW] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com