

# TTF2 Rabbit mAb [BK1K]

Cat NO. :A46906

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	Q9UNY4	130 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

ApplicationDilutionWB1:1000-2000IHC1:100,ICC/IF1:100,The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human TTF2.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

### Subcellular location:

Cytoplasm. Nucleus.

#### Function:

DsDNA-dependent ATPase which acts as a transcription termination factor by coupling ATP hydrolysis with removal of RNA polymerase II from the DNA template. May contribute to mitotic transcription repression. May also be involved in pre-mRNA splicing..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## TTF2 Rabbit mAb [BK1K] Images



Western blot (SDS PAGE) analysis of extracts from Daudi cells lyastes.using TTF2 Rabbit mAb [BK1K] at dilution of 1:1000 incubated at  $4^{\circ}\mathrm{C}$  over night

View more information on http://naturebios.com