

Stanniocalcin 2/STC-2 Rabbit mAb [0798]

Cat NO. :A50043

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	O76061	33/35 kDa	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution
	wв	1:1000-2000
	ІНС	1:100
	ICC/IF	1:100
	The optimal dilutions should be d	etermined by the end user

\sim	•		•
1''	าทแ	100	to.
\mathbf{v}	JIII	uga	LE:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human Stanniocalcin 2/STC-2

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in a variety of tissues including muscle, heart, pancreas, kidney, spleen, prostate, small intestine, colon and peripheral blood leukocytes.

Subcellular location:

Secreted.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Has an anti-hypocalcemic action on calcium and phosphate homeostasis.

Validation Data:

Stanniocalcin 2/STC-2 Rabbit mAb [0798] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells.Using Stanniocalcin 2/STC-2Rabbit mAb [0798] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com