

FXR1 Rabbit mAb[W1S3]

Cat NO. :A50066

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	H,M,R	P51114	70-80KDa	Rabbit	IgG	50ul 100ul,200ul

Applications detail:

Application

WB

1:1000-2000

ICC/IF

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human FXR1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Expressed in all tissues examined including heart, brain, kidney and testis..

Subcellular location:

Cytoplasm, cytosol.

Function:

RNA-binding protein required for embryonic and postnatal development of muscle tissue (PubMed:30770808).

May regulate intracellular transport and local translation of certain mRNAs (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

FXR1 Rabbit mAb[W1S3] Images



Western blot (SDS PAGE) analysis of extracts from 293T cells.Using FXR1 Rabbit mAb IgG [W1S3] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com