

VAMP1 Rabbit mAb[H77T]

Cat NO. :A51553

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M,R	P23763	16kDa	Rabbit	IgG	50ul 100ul,200ul

Applications detail:

Application	Dilution			
WB	1:1000-2000			
ІНС	1:100			
ICC/IF	1:100			
The optimal dilutions should be determined by the end user				

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human VAMP1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Nervous system, skeletal muscle and adipose tissue.

Subcellular location:

[Isoform 1]: Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane, Single-pass type IV membrane protein. Cell junction, synapse, synaptosome., [Isoform 2]: Cytoplasmic vesicle

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Involved in the targeting and/or fusion of transport vesicles to their target membrane.

Validation Data:

VAMP1 Rabbit mAb[H77T] Images



Western blot (SDS PAGE) analysis of extracts from Mouse brain tissue lysate. Using VAMP1 Rabbit mAb IgG [H77T] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com