

FAM98B Rabbit mAb [B5VT]

Cat NO. :A54494

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	Q52LJ0	37 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application Dilution

WB 1:1000-2000

IHC 1:100,
ICC/IF 1:100,
The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human FAM98B.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed strongly in colorectal cancer tissues compared to wild-type colon samples (at protein level)

(PubMed:28040436). Expressed strongly in colorectal cancer tissues compared to wild-type colon

Subcellular location:

Nucleus. Cytoplasm.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Positively stimulates PRMT1-induced protein arginine dimethylated arginine methylation (PubMed:28040436). Promotes colorectal cancer cell malignancy (PubMed:28040436)..

Validation Data:

FAM98B Rabbit mAb [B5VT] Images



Western blot (SDS PAGE) analysis of extracts from Raji cells lyastes.using FAM98B Rabbit mAb [B5VT] at dilution of 1:1000 incubated at 4°C over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.