

# CDC40 Rabbit mAb [78Z3]

Cat NO. :A55734

### Information:

| Applications  | Reactivity: | UniProt ID: | MW(kDa) | Host   | Isotype | Size        |
|---------------|-------------|-------------|---------|--------|---------|-------------|
| WB,IHC,ICC/IF | н           | O60508      | 66 kDa  | Rabbit | IgG     | 100ul,200ul |

Applications detail:

| Application  | Dilution    |  |  |  |
|--|-------------|--|--|--|
| WB   | 1:1000-2000 |  |  |  |
| IHC  | 1:100,      |  |  |  |
| ICC/IF   | 1:100,      |  |  |  |
| The optimal dilutions should be determined by the end user |             |  |  |  |

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human CDC40.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

### Tissue specificity:

### **Subcellular location:**

Nucleus. Nucleus speckle.

#### Function:

Required for pre-mRNA splicing as component of the activated spliceosome (PubMed:33220177). Plays an important role in embryonic brain development, this function does not require proline isomerization (PubMed:33220177)..

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



# **Validation Data:**

# CDC40 Rabbit mAb [78Z3] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells lyastes.using CDC40 Rabbit mAb [78Z3] at dilution of 1:1000 incubated at  $4\,^\circ\mathrm{C}$  over night

View more information on http://naturebios.com