LHFPL5 Rabbit mAb [DMO8]

Cat NO. :A55968

Information:

[Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
	WB	H,M,R	Q8TAF8	24 kDa	Rabbit	lgG	100ul,200ul

Applications detail:

Application Dilution WB 1:1000-2000 The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human LHFPL5.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Cell membrane, Multi-pass membrane protein.

Function:

In the inner ear, may be a component of the hair cell's mechanotransduction machinery that functionally couples PCDH15 to the transduction channel. Regulates transducer channel conductance and is required for fast channel adaptation (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

LHFPL5 Rabbit mAb [DMO8] Images



Western blot (SDS PAGE) analysis of extracts from Human fetal brain tissue lyaste.using LHFPL5 Rabbit mAb [DMO8] at dilution of 1:1000 incubated at 4° over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.

For Research Use Only. Not For Use In Diagnostic Procedures.