

Placental lactogen (CSH1) Rabbit mAb[BF9Q]

Cat NO. :A57570

Information:

| Applications | Reactivity: | UniProt ID: | MW(kDa) | Host | Isotype | Size |
|--------------|-------------|-------------|---------|--------|---------|------------------|
| IHC, ICC/IF | н | P01243 | | Rabbit | IgG | 50ul 100ul,200ul |

| Applications detail: | Application | Dilution | | |
|----------------------|-------------------------------------|--|--|--|
| | IHC | 1:100 | | |
| | ICC/IF | 1:100 | | |
| | The optimal dilutions should be det | The optimal dilutions should be determined by the end user | | |

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human Placental lactogen (CSH1).

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Secreted.

Function:

Produced only during pregnancy and is involved in stimulating lactation, fetal growth and metabolism. Does not interact with GHR but only activates PRLR through zinc-induced dimerization..

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

Placental lactogen (CSH1) Rabbit mAb[BF9Q] Images

Western blot (SDS PAGE) analysis of extracts from .Using Placental lactogen (CSH1) Rabbit mAb IgG [BF9Q] at dilution of 1:1000 incubated at 4°C over night.

View more information on http://naturebios.com