

FRA1 Rabbit mAb [8366]

Cat NO. : A57809

Information:

Nucleus. Function:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	н	P15407	40 kDa	Rabbit	IgG	100ul,200ul

н detail:	P15407 Applicatio	40 kDa	Rabbit	IgG	100ul,200ul		
detail:	Application						
detail:	Application						
	• •	Application Dilution					
	WB				1:1000-2000		
	ICC/IF				1:100,		
	The optimal	The optimal dilutions should be determined by the end user					
on							
ed by immunizi	ng animals with a syr	nthetic peptide o	of Human F	RA1.			
er and cond	litions:						
0 mM PBS, 0.5	mg/ml BSA, 50% glyc	erol (buffer) .					
ore at-20°C or -	80°C.						
for one natural	year of receipt. Avoid	d repeated freez	e / thaw cy	/cles.			
icity:							
	er and cond 0 mM PBS, 0.5 ore at-20°C or -	on sed by immunizing animals with a syner and conditions: 0 mM PBS, 0.5mg/ml BSA, 50% glycore at-20°C or -80°C. for one natural year of receipt. Avoid	on sed by immunizing animals with a synthetic peptide of the conditions: 0 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). ore at-20°C or -80°C. for one natural year of receipt. Avoid repeated freez	on sed by immunizing animals with a synthetic peptide of Human Fer and conditions: 0 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). ore at-20°C or -80°C. for one natural year of receipt. Avoid repeated freeze / thaw cy	on eed by immunizing animals with a synthetic peptide of Human FRA1. er and conditions: 0 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). ore at-20°C or -80°C. for one natural year of receipt. Avoid repeated freeze / thaw cycles.		


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry/ ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey VIr: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

FRA1 Rabbit mAb [8366] Images



Western blot (SDS PAGE) analysis of extracts from HL60 cells lyastes.using FRA1 Rabbit mAb [8366] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night

View more information on http://naturebios.com