ABCF1 Rabbit mAb [LCR2]

Cat NO. :A61732

Information:

ſ	Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
	WB,IHC,ICC/IF	H,R	Q8NE71	95 kDa	Rabbit	lgG	50ul,100ul,200ul

Applications detail:

Application	Dilution		
wв	1:1000-2000		
ІНС	1:100		
ICC/IF	1:100		
The optimal dilutions should be dete	The optimal dilutions should be determined by the end user		

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human ABCF1

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Ubiquitous..

Subcellular location:

[Isoform 2]: Cytoplasm. Nucleus, nucleoplasm. Nucleus envelope.

Function:

Isoform 2 is required for efficient Cap- and IRES-mediated mRNA translation initiation. Isoform 2 is not involved in the ribosome biogenesis..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

ABCF1 Rabbit mAb [LCR2] Images



Western blot(SDS PAGE) analysis of extracts from HeLa cells.Using ABCF1Rabbit mAb [LCR2] at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.

For Research Use Only. Not For Use In Diagnostic Procedures.