

**SLC25A17 Mouse mAb[FMJ9]**

**Cat NO. :A63338**

**Information:**

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M	O43808	35kDa	Mouse	IgG	50ul 100ul,200ul

**Applications detail:**

Application	Dilution
WB	1:1000-2000
IHC	1:100
The optimal dilutions should be determined by the end user	

**Conjugate:**

UnConjugate

**Form:**

Liquid

**sensitivity:**

Endogenous

**Purification:**

Protein A purification

**Specificity:**

Antibody is produced by immunizing animals with a synthetic peptide of human SLC25A17.

**Storage buffer and conditions:**

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

**Tissue specificity:**

Ubiquitous. Expressed in liver..

**Subcellular location:**

Cytoplasm. Peroxisome membrane,Multi-pass membrane protein.

**Function:**

Peroxisomal transporter for multiple cofactors like coenzyme A (CoA), flavin adenine dinucleotide (FAD), flavin mononucleotide (FMN) and nucleotide adenosine monophosphate (AMP), and to a lesser extent for nicotinamide adenine dinucleotide (NAD(+)), adenosine diphosphate (ADP) and adenosine 3',5'-diphosphate (PAP). May catalyze the transport of free CoA, FAD and NAD(+) from the cytosol into the peroxisomal matrix by a counter-exchange mechanism. Inhibited by pyridoxal 5'-phosphate and bathophenanthroline in vitro..

**Introduction:** **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


**Cross Reactivity:** **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Ml:** mink **C:** chicken **Dm** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Hr:** horse

**For Research Use Only. Not For Use In Diagnostic Procedures.**

## Validation Data:

### SLC25A17 Mouse mAb[FMJ9] Images

---



Western blot (SDS PAGE) analysis of extracts from MCF-7 cells. Using SLC25A17 Mouse mAb IgG [FMJ9] at dilution of 1:1000 incubated at 4°C over night.

View more information on <http://naturebios.com>

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.