

TUFM Rabbit mAb [40AS]

Cat NO. :A65583

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	н	P49411	50 kDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:	Application	Dilution	
	WB	1:1000-2000	
	ICC/IF	1:100,	
	The optimal dilutions should be determined by the end user		

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human TUFM.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Mitochondrion.

Function:

Promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis. Plays also a role in the regulation of autophagy and innate immunity. Recruits ATG5-ATG12 and NLRX1 at mitochondria and serves as a checkpoint of the RIG-I/DDX58-MAVS pathway. In turn, inhibits RLR-mediated type I interferon while promoting autophagy..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

TUFM Rabbit mAb [40AS] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells lyastes.using TUFM Rabbit mAb [40AS] at dilution of 1:1000 incubated at $4\,^\circ\!\mathrm{C}$ over night

View more information on http://naturebios.com