

Emi1 Rabbit mAb [M8HL]

Cat NO. :A69751

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H	Q9UKT4	56 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application	Dilution
WB	1:1000-2000
IHC	1:100,
ICC/IF	1:100,
The optimal dilutions should be determined by the end user	

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human Emi1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at -20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, spindle.

Function:

Regulator of APC activity during mitotic and meiotic cell cycle (PubMed:17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:16921029). During mitotic cell cycle plays a role as both substrate and inhibitor of APC-FZR1 complex (PubMed:29875408, PubMed:17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:16921029). During G1 phase, plays a role as substrate of APC-FZR1 complex E3 ligase (PubMed:29875408). Then switches as an inhibitor of APC-FZR1 complex during S and G2 leading to cell-cycle commitment (PubMed:29875408). As APC inhibitor, prevents the degradation of APC substrates at multiple levels: by interacting with APC and blocking

Introduction: **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


Cross Reactivity: **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Ml:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Hr:** horse

For Research Use Only. Not For Use In Diagnostic Procedures.

access of APC substrates to the D-box coreceptor, formed by FZR1 and ANAPC10, by suppressing ubiquitin ligation and chain elongation by APC by preventing the UBE2C and UBE2S activities (PubMed:23708605, PubMed:23708001, PubMed:16921029). Plays a role in genome integrity preservation by coordinating DNA replication with mitosis through APC inhibition in interphase to stabilize CCNA2 and GMNN in order to promote mitosis and prevent rereplication and DNA damage-induced cellular senescence (PubMed:17234884, PubMed:17485488, PubMed:17875940). During oocyte maturation, plays a role in meiosis through inactivation of APC-FZR1 complex. Inhibits APC through RPS6KA2 interaction that increases FBXO5 affinity for CDC20 leading to the metaphase arrest of the second meiotic division before fertilization (By similarity). Controls entry into the first meiotic division through inactivation of APC-FZR1 complex (By similarity). Promotes migration and osteogenic differentiation of mesenchymal stem cells (PubMed:29850565)..

Validation Data:

Emi1 Rabbit mAb [M8HL] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells lysates using Emi1 Rabbit mAb [M8HL] at dilution of 1:1000 incubated at 4°C overnight

View more information on <http://naturebios.com>

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.