

HMGA1 Rabbit mAb[K3LL]

Cat NO. :A70196

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size	
WB,IHC,ICC/IF	H,M,R	P17096	17kDa	Rabbit	IgG	50ul 100ul,200ul	

Applications detail:

Application

WB 1:1000-2000

IHC 1:100

ICC/IF 1:100

The optimal dilutions should be determined by the end user

					te:		
	\sim 10		~	~	-	-	
\mathbf{u}	UH	IIU	u	а	LE	=	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human HMGA1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Nucleus. Chromosome.

Function:

HMG-I/Y bind preferentially to the minor groove of A+T rich regions in double-stranded DNA. It is suggested that these proteins could function in nucleosome phasing and in the 3'-end processing of mRNA transcripts. They are also involved in the transcription regulation of genes containing, or in close proximity to A+T-rich regions.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

HMGA1 Rabbit mAb[K3LL] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells.Using HMGA1 Rabbit mAb lgG [K3LL] at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com