

RhoC Rabbit mAb [2HE0]

Cat NO. :A75780

Information:

| Applications | Reactivity: | UniProt ID: | MW(kDa) | Host | Isotype | Size |
|--------------|-------------|-------------|---------|--------|---------|-------------|
| WB | H,M | P08134 | 21 kDa | Rabbit | IgG | 100ul,200ul |

| Applications detail: | Application | Dilution | |
|----------------------|-----------------------------------|---------------------------|--|
| | wв | 1:1000-2000 | |
| | | | |
| | | | |
| | The optimal dilutions should be d | etermined by the end user | |

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of Human RhoC

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Cell membrane, Lipid-anchor, Cytoplasmic side. Cleavage furrow.

Function:

Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers. Serves as a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Regulates apical junction formation in bronchial epithelial cells..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

RhoC Rabbit mAb [2HE0] Images



Western blot (SDS PAGE) analysis of extracts from U-2 OS cells.Using RhoC Rabbit mAb [2HE0] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com