

# STK39 Rabbit mAb[29F9]

Cat NO. :A77038

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M,R	Q9UEW8	71kDa	Rabbit	IgG	50ul 100ul,200ul

Applications detail:	Application	Dilution		
	WB	1:1000-2000		
	IHC	1:100		
	ICC/IF	1:100		
	The optimal dilutions should be d	The optimal dilutions should be determined by the end user		

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human STK39.

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

### Tissue specificity:

Predominantly expressed in brain and pancreas followed by heart, lung, kidney, skeletal muscle, liver, placenta and testis.

## Subcellular location:

Cytoplasm. Nucleus.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



May act as a mediator of stress-activated signals. Mediates the inhibition of SLC4A4, SLC26A6 as well as CFTR activities by the WNK scaffolds, probably through phosphorylation. Phosphorylates RELT..

# **Validation Data:**

### STK39 Rabbit mAb[29F9] Images



Western blot (SDS PAGE) analysis of extracts from Mouse testis tissue lysate. Using STK39 Rabbit mAb IgG [29F9] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com