

# p47-phox Rabbit mAb [R65I]

Cat NO. :A77299

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	н	P14598	45 kDa	Rabbit	IgG	100ul,200ul

Applications detail: Application Dilution

WB 1:1000-2000

IHC 1:100,

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human p47-phox.

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

Detected in peripheral blood monocytes and neutrophils (at protein level)..

## Subcellular location:

Cytoplasm, cytosol. Membrane, Peripheral membrane protein, Cytoplasmic side.

#### Function:

NCF2, NCF1, and a membrane bound cytochrome b558 are required for activation of the latent NADPH oxidase (necessary for superoxide production)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



### **Validation Data:**

### p47-phox Rabbit mAb [R65I] Images



Western blot (SDS PAGE) analysis of extracts from Raji cells lyastes.using p47-phox Rabbit mAb [R65I] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night

View more information on http://naturebios.com