

EPHX2 Rabbit mAb [2762]

Cat NO. :A78625

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H	P34913	63 kDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:

Application	Dilution
WB	1:1000-2000
The optimal dilutions should be determined by the end user	

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human EPHX2.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at -20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Cytoplasm. Peroxisome.

Function:

Bifunctional enzyme (PubMed:12574510). The C-terminal domain has epoxide hydrolase activity and acts on epoxides (alkene oxides, oxiranes) and arene oxides (PubMed:12869654, PubMed:12574510, PubMed:22798687). Plays a role in xenobiotic metabolism by degrading potentially toxic epoxides (By similarity). Also determines steady-state levels of physiological mediators (PubMed:12869654, PubMed:12574510, PubMed:22798687, PubMed:21217101)... Bifunctional enzyme (PubMed:12574510). The N-terminal domain has lipid phosphatase activity, with the highest activity towards threo-9,10-phosphonoxy-hydroxy-octadecanoic acid, followed by erythro-9,10-phosphonoxy-hydroxy-octadecanoic acid, 12-phosphonoxy-octadec-9Z-enoic acid and 12-

Introduction: **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


Cross Reactivity: **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Ml:** mink **C:** chicken **Dm** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Hr:** horse

For Research Use Only. Not For Use In Diagnostic Procedures.

phosphonoxy-octadec-9E-enoic acid (PubMed:12574510). Has phosphatase activity toward lyso-glycerophospholipids with also some lower activity toward lysolipids of sphingolipid and isoprenoid phosphates (PubMed:22217705, PubMed:22387545)..

Validation Data:

EPHX2 Rabbit mAb [2762] Images



Western blot (SDS PAGE) analysis of extracts from 293T cells lysates using EPHX2 Rabbit mAb [2762] at dilution of 1:1000 incubated at 4°C over night

View more information on <http://naturebios.com>

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.