smooth muscle Myosin heavy chain 11 Rabbit mAb [YNAE]

Cat NO. :A79600

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	н	P35749	227 kDa	Rabbit	lgG	100ul,200ul

Applications detail:

Application	Dilution
WB	1:1000-2000
IHC	1:100
The optimal dilutions should b	e determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human smooth muscle Myosin heavy chain 11

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Smooth muscle, expressed in the umbilical artery, bladder, esophagus and trachea. Isoform 1 is mostly found in

slowly contracting tonic muscles..

Subcellular location:

Melanosome.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Muscle contraction.

Validation Data:

smooth muscle Myosin heavy chain 11 Rabbit mAb [YNAE] Images



Western blot (SDS PAGE) analysis of extracts from Human bladder.Using smooth muscle Myosin heavy chain 11Rabbit mAb [YNAE] at dilution of 1:1000 incubated at

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.