

HNRNPA0 Rabbit mAb [NZ4C]

Cat NO. :A81007

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M,R	Q13151	32 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application Dilution

WB 1:1000-2000

IHC 1:100

ICC/IF 1:100

The optimal dilutions should be determined by the end user

		gate:			
1 · ^	nıı	-	to.		
-		. 9 4			

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human HNRNPA0

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Nucleus.

Function:

mRNA-binding component of ribonucleosomes. Specifically binds AU-rich element (ARE)-containing mRNAs. Involved in post-transcriptional regulation of cytokines mRNAs..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

HNRNPA0 Rabbit mAb [NZ4C] Images



Western blot (SDS PAGE) analysis of extracts from HEK293T cells.Using HNRNPA0Rabbit mAb [NZ4C] at dilution of 1:1000 incubated at $4\,^\circ\!\mathrm{C}$ over night.

View more information on http://naturebios.com